

Proceedings of the Iowa Academy of Science

Volume 29 | Annual Issue

Article 24

1922

Classification of Lenses

Louise Fillman
State University of Iowa

Copyright © Copyright 1922 by the Iowa Academy of Science, Inc.
Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Fillman, Louise (1922) "Classification of Lenses," *Proceedings of the Iowa Academy of Science*, 29(1), 128-128.
Available at: <https://scholarworks.uni.edu/pias/vol29/iss1/24>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

of a very interesting Eurypterid. The carapace is sub-semicircular in outline; anterior margin well rounded; posterior margin broadly concave; sides diverging gently posteriorly and somewhat produced at the postero-lateral angles. The compound eyes are prominent, bean-shaped, situated about in the middle, and as far apart as their distance from the outer edge. Length, 8 mm., width, 13 mm. The species is named *Eurypterus thomasi* in honor of Prof. A. O. Thomas.

DEPARTMENT OF GEOLOGY,
STATE UNIVERSITY OF IOWA.

ORIGIN OF LIMESTONE CONGLOMERATES

LOUISE FILLMAN

A study of the literature reveals at least eight ways in which limestone conglomerates may be formed. The following ways are noted and described in the paper: (1) derivatives from some pre-existing rock; (2) by deformation of the laminæ; (3) by pebbles derived from concretions; (4) by the secretions of limestone nodules by algae; (5) by the breaking of thin layers of limestone by storm waves; (6) by subaquatic gliding of limestone layers; (7) by lime-mud being cracked on a tidal beach; (8) by minor oscillations of the sea and the erosion of materials previously deposited.

CLASSIFICATION OF LENSES

LOUISE FILLMAN

This paper is the result of a bibliographic study of lenses, on the part of Lloyd North and the writer. In the literature some thirty-two genetic types are described. In the present paper these types are classified and described. A bibliography is appended.

DEPARTMENT OF GEOLOGY,
STATE UNIVERSITY OF IOWA.

THE ROCKFORD GEODES

S. L. GALPIN

These geodes occur in Lime Creek shales. They are rather unusual in containing a number of roundish cavities rather than the customary rough opening. The cavities are lined with: 1, small